

International Application No. PCT/EP98/01117
 Date: August 23, 1999

Attorney Docket No. DECLE2.001APC

09/380106 Page 1

**TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US)
 CONCERNING A FILING UNDER 35 USC 371**

International Application No.: PCT/EP98/01117
 International Filing Date: 19 February 1998
 Priority Date Claimed: 21 February 1997
 Title of Invention: PLANT MICROPROPAGATION AND GERMPLASM STORAGE
 Applicant(s) for DO/EO/US: Koenraad Hilaire Maria GOETHALS; Mondher EL JAZIRI; Marc Charles Ernest VAN MONTAGU

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. (X) This is a **FIRST** submission of items concerning a filing under 35 USC 371.
2. (X) This express request to begin national examination procedures (35 USC 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 USC 371(b) and PCT Articles 22 and 39(1).
3. (X) A proper Demand for International Preliminary Examination was made by the 19th month from the earliest claimed priority date.
4. (X) A copy of the International Application as filed (35 USC 371(c)(2)) has been transmitted by the International Bureau.
5. (X) Amendments to the claims of the International Application under PCT Article 19 (35 USC 371(c)(3)) have not been made and will not be made.
6. (X) A copy of the International Preliminary Examination Report with any annexes thereto, such as any amendments made under PCT Article 34.
7. (X) A **FIRST** preliminary amendment.
8. (X) International Application as published.
9. (X) A return prepaid postcard.
10. (X) The filing fees are calculated below:

BASIC FEE			\$840	
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE	
Total Claims	10 - 20 =	0 ×	\$18	\$0
Independent Claims	1 - 3 =	0 ×	\$78	\$0
Multiple dependent claims(s) (if applicable)			\$260	\$0
TOTAL OF ABOVE CALCULATIONS			\$840	

11. (X) A check in the amount of \$840 to cover the above fees is enclosed.

-
12. (X) The fee for later submission of the signed oath or declaration set forth in 37 CFR 1.492(e) will be paid upon submission of the declaration.
13. (X) The Commissioner is hereby authorized to charge only those additional fees which may be required to avoid abandonment of the application, or credit any overpayment to Deposit Account No. 11-1410. A duplicate copy of this sheet is enclosed.

SEND ALL CORRESPONDENCE TO:

KNOBBE, MARTENS, OLSON & BEAR, LLP
(Customer No. 20,995)



Daniel E. Altman
Registration No. 34,115

E:\DOCS\DEA\DEA-1572.DOC:kc
082399

664447" 90708660

VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL-ENTITY STATUS

I, the undersigned, do hereby declare that:

☒ I am an official empowered to act on behalf of the nonprofit organization identified below:

NAME OF ORGANIZATION: Vlaams Interuniversitair Instituut Voor Biotechnologie

ADDRESS OF ORGANIZATION: Rijvisschestraat 118, Bus 1, D-9052 Zwijnaarde, Belgium

TYPE OF ORGANIZATION:

NAME OF ORGANIZATION: Universite' Libre de Bruxelles

ADDRESS OF ORGANIZATION: Avenue F.D. roosevelt 50, B. 1050 Brussels, Belgium

TYPE OF ORGANIZATION:

☒ University or other institution of higher education.

I further declare that the nonprofit organization identified above qualifies as a nonprofit organization as defined in 37 CFR 1.9(e) for purposes of paying reduced fees to the United States Patent and Trademark Office regarding the invention described in the patent or application identified above.

The individual, concern or organization identified above has not assigned, granted, conveyed or licensed, and is under no obligation under contract or law to assign, grant, convey or license, any rights in the invention to any person who would not qualify as an independent inventor under 37 CFR 1.9(c) if that person had made the invention, or to any concern which would not qualify as a small business concern under 37 CFR 1.9(d) or a nonprofit organization under 37 CFR 1.9(e).

If the rights held by the above-identified individual, concern or organization are not exclusive, each individual, concern or organization having rights in the invention are identified below. Each such individual, concern or organization must file separate verified statements averring to their status as small entities.

***NOTE:** Separate verified statements are required from each named person, concern or organization having rights to the invention averring to their status as small entities. (37 CFR 1.27).

FULL NAME: Vlaams Interuniversitair Instituut voor Biotechnologie

ADDRESS: Rijvisschestraat 120, B-9052 Zwijnaarde, Belgium

☐ INDIVIDUAL ☐ SMALL BUSINESS CONCERN ☐ NONPROFIT ORGANIZATION

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small-entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 CFR 1.28(b)).

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to which this verified statement is directed.

VERIFIED STATEMENT (DECLARATION) CLAIMING SMALL-ENTITY STATUS

I, the undersigned, do hereby declare that:

☒ I am an official empowered to act on behalf of the nonprofit organization identified below:

NAME OF ORGANIZATION: Vlaams Interuniversitair Instituut Voor Biotechnologie

ADDRESS OF ORGANIZATION: Rijvisschestraat 118, Bus 1, D-9052 Zwijnaarde, Belgium

TYPE OF ORGANIZATION:

NAME OF ORGANIZATION: Universite' Libre de Bruxelles

ADDRESS OF ORGANIZATION: Avenue F.D. roosevelt 50, B. 1050 Brussels, Belgium

TYPE OF ORGANIZATION:

☒ University or other institution of higher education.

I further declare that the nonprofit organization identified above qualifies as a nonprofit organization as defined in 37 CFR 1.9(e) for purposes of paying reduced fees to the United States Patent and Trademark Office regarding the invention described in the patent or application identified above.

The individual, concern or organization identified above has not assigned, granted, conveyed or licensed, and is under no obligation under contract or law to assign, grant, convey or license, any rights in the invention to any person who would not qualify as an independent inventor under 37 CFR 1.9(c) if that person had made the invention, or to any concern which would not qualify as a small business concern under 37 CFR 1.9(d) or a nonprofit organization under 37 CFR 1.9(e).

If the rights held by the above-identified individual, concern or organization are not exclusive, each individual, concern or organization having rights in the invention are identified below. Each such individual, concern or organization must file separate verified statements averring to their status as small entities.

***NOTE:** Separate verified statements are required from each named person, concern or organization having rights to the invention averring to their status as small entities. (37 CFR 1.27).

FULL NAME: Universite' Libre de Bruxelles

ADDRESS: Avenue F.D. Roosevelt 50, B-1050 Brussels, Belgium

☐ INDIVIDUAL ☐ SMALL BUSINESS CONCERN ☐ NONPROFIT ORGANIZATION

I acknowledge the duty to file, in this application or patent, notification of any change in status resulting in loss of entitlement to small-entity status prior to paying, or at the time of paying, the earliest of the issue fee or any maintenance fee due after the date on which status as a small entity is no longer appropriate. (37 CFR 1.28(b)).

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application, any patent issuing thereon, or any patent to which this verified statement is directed.

NAME OF PERSON SIGNING: Rudy Dekeyser

TITLE OF PERSON (if not an owner or individual): Vice General Director & Technology Transfer Manager

ADDRESS OF PERSON SIGNING: Rijvisschestraat 118, Bus 1, D-9052 Zwijnaarde, Belgium

SIGNATURE: 

DATE: 22/10/99

NAME OF PERSON SIGNING: Jean-Louis Vanherweghem


TITLE OF PERSON (if not an owner or individual): Recteur

ADDRESS OF PERSON SIGNING: Avenue F.D. Roosevelt 50, B. 1050 Brussels, Belgium

SIGNATURE: 

DATE: 18/10/1999

DEA-1565.DOCsIs
092899



66421 " 90F08E60

09/380106

DECL2.001APC

PATENT

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant	:	Goethals, et al.)	Group Art Unit Unknown
)	
Int'l)	
Appl. No.	:	PCT/EP98/01117)	
)	
Int'l Filing)	
Date	:	February 19, 1998)	
)	
For	:	PLANT MICROPROPAGATION)	
		AND GERMPLASM STORAGE)	
)	
Examiner	:	Unknown)	
)	

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents
Washington, D.C. 20231

Dear Sir:

Preliminary to examination on the merits, please amend the above-captioned U.S. National Phase Application as follows:

IN THE SPECIFICATION:

On page 1 of the Specification, after the Title of the Invention and before the Field of the Invention statement starting on line 1, please insert --This is the U.S. National Phase under 35 U.S.C. § 371 of International Application PCT/EP98/01117, filed February 19, 1998.--.

On page 16 before Claim 1, please delete the word "CLAIMS" and substitute therefor --WHAT IS CLAIMED IS--.

IN THE CLAIMS:

1. (Amended) [Method]A method for [the *in vitro* preparation of starting materials for]plant micropropagation and/or storage of germplasm, comprising the steps of:

(a) contacting plant material with a [**micro-organism**]microorganism that induces fasciation and/or one or more fasciation-inducing factors derived from the [**micro-organism**]microorganism or derived from the infected and/or fasciated tissue;

(b) developing leafy galls or shoot outgrowths on the plant material; and

(c) isolating the leafy galls or shoot outgrows as the starting materials.

2. (Amended) [Method]The method for [the *in vitro* micropropagation of plants, which method comprises the method as claimed in] plant micropropagation and/or storage of germplasm of claim 1, further [characterized by] comprising the steps of:

(d) eliminating or inactivating the [micro-organism]microorganism that induces fasciation and/or the one or more fasciation-related factors derived from the [micro-organism]microorganism or derived from the infected and/or fasciated tissue;

(e) culturing the leafy gall or shoot outgrows in or on one or more suitable culture media to allow shoot and root development for obtaining plantlets; and

(f) transferring the plantlets thus obtained, [optionally after acclimatisation,] to conventional growing conditions to obtain regenerated plants.

3. (Amended) [Method]The method for [the preservation of plants or germplasm, which method comprises the method as claimed in] plant micropropagation and/or storage of germplasm of claim 1, further [characterized by]comprising the step of storing the leafy galls or shoot outgrows under growth limiting conditions.

4. (Amended) [Method]The method [as claimed in claims 1-3]of claim 1, wherein the fasciation-inducing [micro-organism]microorganism is *Rhodococcus fascians*.

5. (Amended) [Method]The method [as claimed in claim 3 and 4]of claim 3, wherein the growth limiting conditions comprise [low] temperatures of about 4 °C or less or [cryo-conserving conditions, such as]or wherein the growth limiting conditions comprise storage in liquid nitrogen.

6. (Amended) [Method]The method [as claimed in claims 1-5]of claim 1, wherein the plant material [comprises explants, such as]is selected from the group consisting of: leaf discs, organs, tissue fragments, seedlings, seeds, embryos, isolated cells, protoplasts, cell cultures and callus tissues.

7. (Amended) [Method]The method [as claimed in claims 1-6]of claim 1, wherein the plant material is material of species of the genera selected from the group consisting of: Brassicaceae, Salicaceae, Compositae, Solanaceae, Scrophulariaceae, Liliaceae, Papaveraceae, Gramineae and Fabaceae.

8. (Amended) [Method]The method for the [in vitro micropropagation of plants, which method comprises the method of] plant micropropagation and/or storage of germplasm of claim 3 [and optionally]further comprising: [the subsequent]

remov[all]ing or inactivat[ion]ing [of] microorganisms still present [micro-organisms] that induce fasciation and/or removing or inactivating fasciation related factors derived from the [micro-organism]microorganism or derived from the infected and/or fasciated tissue,

culturing the leafy gall or shoot outgrows in or on one or more suitable culture media to induce shoot and root development for obtaining plantlets, and

transferring the plantlets thus obtained, [optionally after acclimatisation,] to conventional growing conditions to obtain regenerated plants.

Please add the following Claims:

9. The method of claim 2 wherein the plantlets are acclimatized before conventional growing conditions to obtain regenerated plants.

10. The method of claim 8 wherein the plantlets are acclimatized before conventional growing conditions to obtain regenerated plants.

REMARKS

The claims have been amended to conform with the rules of practice before the Patent and Trademark Office. The specification has been amended to recite the International Application and priority application. The word "CLAIMS" has been deleted and substituted by "WHAT IS CLAIMED IS" so that subsequently appearing claims will be the object of a sentence as specified by MPEP § 608.01(m). As a result of this preliminary amendment, Claims 9-10 have been added and Claims 1-8 have been amended. Accordingly, Claims 1-10 are presented for examination. No new matter is being added herewith.

Should there be any questions concerning this application, the Examiner is invited to contact the undersigned attorney at the telephone number appearing below.

Respectfully submitted,

KNOBBE, MARTENS, OLSON & BEAR, LLP

Dated:

23 Aug. 1999

By:

Daniel E. Altman

Daniel E. Altman
Registration No. 34,115
Attorney of Record
620 Newport Center Drive
Sixteenth Floor
Newport Beach, CA 92660
(949) 760-0404

426 Rec'd PCT/PTO 23 AUG 1999

09/380106

PCT/EP98/01117

WO 98/36635

KPTS

PLANT MICROPROPAGATION AND GERMPLASM STORAGE

The present invention relates to new methods of plant micropropagation and germplasm storage.

Plants can be propagated through their two developmental life cycles: the sexual or the asexual (or vegetative). In the vegetative (asexual) cycle the unique characteristics of any individual plant selected for propagation are usually perpetuated. In most cases, each new plant produced by this method may be considered to be an extension of the somatic cell line of one individual.

10 Methods available for propagating plants in vitro (micro-propagation) are largely an extension of those already developed for conventional propagation (macro-propagation).

The methods that are theoretically available for propagating plants in vitro are essentially based on the multiplication of shoots from axillary buds, on the formation of adventitious shoots and on the formation of somatic embryos. These processes are initiated either directly from the original explant or indirectly from unorganised cells such as in callus or cell suspension cultures.

In practice most micropropagated plants are produced at present by the multiplication of shoots from axillary buds. Nevertheless, somatic embryos, shoots and/or plantlets do not always originate in a single way. For example, in shoot cultures, besides axillary shoots, there are sometimes adventitious shoots formed directly on existing leave or stem explants and/or shoots arising indirectly from callus at the base of the explant.

30 Considering these observations, somaclonal variation is frequently observed among the regenerated plants even through multiplication of shoots from axillary buds which is considered as the most stable source for plant micropropagation. In such mixed cultures, shoots arising indirectly from callus are not desirable because of the somaclonal variation. This is particularly important when the method is applied in a large-scale programme (industrial scale). In order to

09380106-12499

WO 98/36635

PCT/EP98/01117

2

reduce this genetic/ epigenetic variation,
standardization of the cellular origin for the
regenerated new shoots is therefore essential.

A second limitation of the method of axillary
5 shoot production is associated with the use of exogenous
phytohormones for plant micro-propagation purposes. The
growth and proliferation of axillary shoots in shoot
cultures is usually promoted by incorporating growth
regulators into the culture medium. Depending on the
10 plant species, usually cytokinins and auxins in
appropriate combinations and concentrations are used.
Such treatments effectively remove the dominance of
apical meristems so that axillary shoots are produced.
These shoots are used as miniature cuttings for plant
15 multiplication.

The cytokinin growth regulators added to shoot
culture media to promote axillary shoot growth, usually
inhibit root formation. Therefore, single shoots must be
moved to a different medium for rooting in vitro
20 (generally medium containing auxins) before being
transferred as juvenile plantlets to the external
environment. The axillary shoot production thus requires
a sequence of steps.

Nowadays, there is an increased tendency to
25 consider that exogenous applied synthetic cytokinin,
which is generally too high, may cause many abnormal
features (side effects) such as production of small
shoots which typically fail to elongate, induction of
unusual leave shapes and particularly the tendency of the
30 regenerated shoots to become vitrified.

In addition, high concentrations of usual
cytokinins generally inhibit or considerably delay root
formation and also prevent root growth and the promoting
effects of auxins on root initiation. In this respect,
35 frequently more than one subculture to a cytokinin-free
medium may be required until the level of cytokinin
within the tissues has been sufficiently reduced followed
by a transfer of the shoots to a rooting medium.

504495 44495

WO 98/36635

PCT/EP98/01117

3

Considering these aspects it is furthermore impossible to propagate particular plant species which are sensitive to such hormonal treatments.

It is thus the object of the present invention to
5 provide an alternative micropropagation method with which the above identified drawbacks of the known methods are obviated.

According to the invention it has surprisingly been found that leafy galls, which are produced on
10 dicotyledonous plants by infection with particular micro-organisms or by treatment with substances produced by these micro-organisms or derived from the infected and/or fasciated tissue (so-called fasciation), and which consist of multiple meristematic centers and shoot
15 primordia that are suppressed for elongation, are an excellent starting point for micropropagation. In a gall no apical dominance phenomena are evident and the formation of numerous primordia is derepressed by the infection process. The gall is in fact a large mass of
20 shoot primordia that are covered with leaves and that are infected by the micro-organism. In leafy galls specific metabolites are present that are involved in the establishment and the maintenance of the leafy gall structure.

25 In monocotyledonous plants the structures that arise upon contact with or infection by fasciation-inducing micro-organisms or substances have a different morphology than the above described leafy galls that are formed in dicotyledonous plant material. Infection either
30 results in deformation of bulbs and/or the formation of several elongated shoots. These monocot elongated shoot structures are useful in the method of the invention and will be further identified as "shoot outgrows", whereas the dicot structures are called "leafy galls".

35 Apart from the morphogenetic aspects, a gall represents a center of elongation inhibition where the bacterium suppresses the outgrowth of the primordia in the gall. This phenomenon is translated in the apparent

0380405-134465

WO 98/36635

PCT/EP98/01117

4

lack of apical dominance that would be exerted by one of the primordia and would lead to the outgrowth of distinct shoot primordia from the gall tissues. As such, each gall can be considered as a highly condensed plant where the amplified production of primordia and meristems -apical as well as axillary- are confined to one compact region through the suppression of both the elongation processes and apical dominance.

Furthermore, it was found according to the invention that due to the fact that the galls consist of large numbers of shoot primordia confined to a compact and small structure, they are also a very suitable material for germplasm storage.

Starting from these two observations the present invention now provides for a method for the in vitro preparation of starting materials for micropropagation and/or storage of germplasm, comprising the steps of:

- a) contacting plant material with a micro-organism that induces fasciation and/or one or more fasciation-inducing factors derived from the micro-organism, or derived from the infected and/or fasciated tissue;
- b) developing leafy galls or shoot outgrows on the plant material; and
- c) isolating the leafy galls or shoot outgrows as the starting materials.

This method of the invention may then be incorporated in a method of micropropagation or a method of germplasm storage, respectively. A method for the in vitro micropropagation of plants comprises the above described method for the preparation of starting material, further characterized by the steps of

- d) eliminating or inactivating the micro-organism that induces fasciation and/or the one or more fasciation-related factors derived from the micro-organism or derived from the infected and/or fasciated tissue;

09380106-113499

WO 98/36635

PCT/EP98/01117

5

e) culturing the leafy gall or shoot outgrows in or on one or more suitable culture media to allow further shoot and root development for obtaining plantlets, and

- 5 f) transferring the plantlets thus obtained, optionally after acclimatisation, to conventional growing conditions to obtain regenerated plants.

A method for the preservation of plants or germplasm comprises the method for the preparation of
10 starting material, further characterized by the step of storing the leafy galls or shoot outgrows under growth limiting conditions. The growth limiting conditions may comprise low temperatures of about 4°C or cryo-conserving conditions, such as storage in liquid nitrogen.

- 15 Both methods are not only applicable on angiosperms, but also on gymnosperms.

In a preferred embodiment of the invention the fasciation inducing micro-organism is the Gram positive bacterium Rhodococcus fascians (Tilford) (Goodfellow M. &
20 Cross, T., "Classification", published: Goodfellow, Modarski & Williams ed., The Biology of Actinomycetes, Academic Press London, p. 7-64 (1984)). (Effective publication Goodfellow M., "Reclassification of Corynebacterium fascians (Tilford, Dowson) in the genus
25 Rhodococcus, as Rhodococcus fasc. comb. nov." System. Appl. Microbiol. 5, 225-229 (1984), previously known as Phytomonas fascians Tilford, J., Agric. Res. 53, 383-394 (1936); or Corynebacterium fascians (Tilford) Dowson, W.J., Trans. Brit. Myc. Soc. 25, 311-314 (1942)).
30 Rhodococcus fascians is a Nocardioform Actinomycete that infects a whole range of both dicotyledonous and monocotyledonous plants.

The range of plants that can be infected by R. fascians is very broad with different species belonging
35 to numerous plant families, such as Brassicaceae, Salicaceae, Compositae, Solanaceae, Scrophulariaceae, Liliaceae, Papaveraceae, Gramineae and Fabaceae.

09380106 112499

WO 98/36635

PCT/EP98/01117

6

Infection of an intact plant can for example be achieved by vacuum infiltration of plants with suspensions of the bacteria, application of a dense bacterial culture sample to a wound surface or, less efficiently, by simply applying bacteria in either the growth substrate of the plants or on the surface of intact plants. Also explants, such as leaf discs, organs, tissue fragments, and seedlings, seeds, embryos, isolated cells and protoplasts can be successfully infected, giving rise to leafy galls or shoot outgrows. Furthermore, R. fascians infected cell cultures and callus tissues react in a similar fashion.

The role of R. fascians in the phenomenon of suppression of elongation and apical dominance can be exemplified by killing of the bacteria in the gall through a treatment with bactericidal agents such as Carbenicillin, Triacellin and Claforan. Growth of detached galls on hormone free plant media with the addition of bactericidal compounds to an end concentration of 500 mg/L, results in the release of the amplified primordia from the bacterial suppression. ensuing in the outgrowth of numerous independent shoots from the gall tissues that will produce root tissues. The rooting can be facilitated by treating the galls in the dark where the release of the shoots is accompanied by etiolation, producing masses of elongated sprouts. Each shoot can be transferred to another medium resulting in the production of a fertile plant clone.

These new micropropagation and germplasm storage methods are widely applicable and may for example be used for trees, crops, medicinal and aromatic plants. The methods are also suitable for biochemical or genetic selection purposes.

Instead of using the micro-organism as a whole one or more factors derived from the micro-organism or infected and/or fasciated tissue may be used. These factors are capable of inducing the formation of leafy galls in dicotyledonous plants and shoot outgrows in

09380406-14499

monocotyledonous plants. The factors may for example be gene products of the genes involved in fasciation, or the molecules produced by these gene products.

In the R. fascians strain D188, the property to induce fasciation is correlated with the presence of a +200 kb large linear plasmid, pFiD188, that carries several loci (e.g. fas, att and hyp) involved in leafy gall formation. The present inventors have found that mutations in the fas locus render the bacteria non-pathogenic. A mutation in the att locus provokes attenuated fasciation resulting in smaller leafy galls that are formed more slowly. Finally, a mutation in the hyp locus results in hyper-fasciation, leading to the formation of larger galls compared to wild type.

Apart from pFiD188 encoded genes, also chromosomal genes in the bacterium are involved in the infection process for instance by influencing the endophytic/epiphytic survival of R. fascians.

Evidently, gall formation results from the modulation of the regular hormone balance in the infected plant tissues. Although the fasciation symptoms superficially resemble effects caused by the application of different doses of several cytokinins, they are nevertheless clearly discriminated from the latter by the presence of non-elongating primordia and the absence of abundantly developing shoots. Early reports where galling was correlated with the presence of different cytokinins in the growth medium of R. fascians have recently been questioned by opposing results. This is further substantiated by the present inventors by their finding of the presence of cytokinins in the growth media of a non-fasciating strain D188-5 (Crespi et al., EMBO J. 11, 795-804 (1992)) cured for pFiD188.

Although early results hinted at the presence of an adenosine monophosphate isopentenyl transferase (ipt) homologous gene in the fas locus on pFiD188 that would be involved in the biosynthesis of cytokinins (Crespi et al., 1992, supra), the present results question the

66444-1-100000

WO 98/36635

PCT/EP98/01117

8

unique involvement of the ipt homologue in the biosynthesis of regular cytokinins like isopentenyl adenine or zeatin. Instead, the fas locus is involved in the production and secretion of one or more isopentenylated molecules that affect the plants hormone levels. This was confirmed by the isolation and preliminary chemical analysis of one fas dependent products and fits with the overall low amino acid identity between the fas encoded ipt gene product and regular isopentenyl transferase proteins. Apart from the ipt related gene, the fas locus encodes several genes that, on the basis of sequence homology studies, are implicated in cytochrome P450 dependent oxygenation and S-Adenosyl Methionine dependent methylation processes of, conceivably, the fas product.

High expression of the fas locus in the bacterium -as measured by translational GUS fusion assays- is strictly controlled and can only be consistently achieved in batch cultures using a defused induction medium, such as Minimal A-medium (obtainable by adding 20.5 ml 0.1M citrate and 29.5 ml 0.1M Na-citrate to 450 ml distilled water and adding after autoclaving and cooling 500 μ l 1% (w/v) thiamine, 1.25 ml MgSO₄ and 1.25 ml 40% NH₄SO₄) and only in the presence of either gall extracts or histidine in combination with particular carbon sources such as pyruvate and succinate. Plant extracts cannot induce fas gene expression, but gall extracts contain a specific inducer molecule. This inducer molecule is present in all gall tissues formed on a whole range of different plants and is able to induce expression in minimal concentrations. Furthermore, production of this molecule was shown in tobacco BY-2 cell cultures confronted with R. fascians wild type cells while this was not the case for non-infected plant cell cultures or pure bacterial cultures.

The fas expression is mainly regulated at the post-transcriptional level, indirectly through the activity of the fasR gene that encodes an AraC type

09380105 1.12499

WO 98/36635

PCT/EP98/01117

9

transcriptional activator and possibly directly through the hyp locus that carries a gene encoding an RNA-helicase.

Confrontation of BY-2 tobacco cell culture
5 synchronized for mitosis using aphidicolin with fas
induced D188 cells resulted in a marked increase of the
number of tobacco cells that enter the first round of
mitosis upon relieve of the mitosis block. This effect is
absent when the tobacco cell culture is confronted with
10 the non-fasciating strain D188-5 or with fas mutant
strains.

Classical cytokinin-bioassays using extracts of
R. fascians D188 and D188-5 grown for fas induction
proved difficult to interpret due to the presence of high
15 background levels of contaminating cytokinins in both
tested extracts. From D188 cultures, induced for fas
expression, a fas dependent metabolite was isolated.

The att locus comprises a 20 kb region of
pFiD188, linked to the fas locus. Sequencing showed the
20 presence of several genes that are involved in the
synthesis and secretion of a complex molecule of unknown
structure. Apart from genes involved in the synthesis of
arginine, the locus carries genes implicated in the
biosynthesis and modification of fatty acid related
25 molecules and a gene related to export protein genes.

Expression of the att locus is partially co-
regulated with the fas locus; induction of GUS fusions is
only visible in defined media in the presence of gall
extracts or of histidine in combination with particular
30 carbon sources and depends on the activity of the fasR
gene. However, an independent regulation through an att
specific regulatory gene, attR, is postulated here
whereby the expression of the attR gene is itself
enhanced by the fasR gene in response to plant extracts.

35 Using ¹⁴C-labelled arginine or acetate, in either
case the induced accumulation in the growth medium of R.
fascians D188 of the same ¹⁴C-labelled, att dependent
compounds was evident using thin layer chromatography,

05330105 "412499

WO 98/36635

PCT/EP98/01117

10

proving the incorporation of both tested substrates in complex, secreted compounds. These were isolated through High Pressure Liquid Chromatography for chemical analysis and bio-assaying using BY-2 tobacco cell cultures. As an example, a molecule with a molecular weight of 681 dalton was isolated.

The products of the att locus are probably involved in the formation of leafy galls in combination with the fas locus as a mutant in the locus is strongly retarded in the galling process. It is also conceivable that the att and fas loci cooperate in the biosynthesis and secretion of a single type of molecules that provoke fasciation.

Apart from the fas and att loci, still other loci on pFid188 are involved in the formation of leafy galls. The hyp locus probably negatively controls expression of the fas and att expression through RNA helicase activity on the mRNA. Further downstream of the fas locus another locus has been isolated that is essential for fasciation and sequencing of which showed the presence of oligopeptide synthetic genes.

Molecular analysis of the gene expression in infected tobacco plants using differential display techniques showed the induction of different genes showing homology to genes involved in the metabolism of gibberellins and steroids as well as the repression of a gene showing homology to oligopeptide receptor proteins. These results strongly suggest an influence of the infecting bacteria and their products on the existing balances of different plant growth factors, resulting in the induction of meristem and primordium formation.

The formation of a leafy gall by R. fascians is thus a complex process with different levels of signalling between both partners. The expression of the bacterial fasciation genes located on the linear plasmid is prone to a complex regulation that ultimately implies the involvement of a gall specific metabolite in the fasR dependent induction of the att and fas genes. Once the

0360106-12499

WO 98/36635

PCT/EP98/01117

11

fas and att genes are optimally expressed, metabolites are produced and secreted in the plant that disturb and/or redirect the hormone household of the plant tissues. The fas and att derived factor(s) are implicated in the enhancement of plant cell divisions and are crucial for the activation of the gall formation. Molecular data suggest the involvement of gibberellins, steroid type molecules and/or possibly oligopeptides of the plant. This unique response of the plant to infection confers the particular properties of R. fascians and its derived products in altering the developmental programming of the plant in some hitherto non-described way. It further discriminates the leafy gall syndrome from other neoplastic pathologies like the Agrobacterium tumefaciens induced crown gall formation and the Pseudomonas savastanoi induced galling of Oleander species.

In summary, it has now been found that a gall constitutes several meristems and primordia the formation of which is correlated with the fas product, the att products, or a combination of both and possibly other products the synthesis of which is determined by different loci on pFiD188 or on the bacterial chromosome. The end result is a local activation and deregulation of primordium and meristem formation that results in dense tissues -the gall- that is ultimately covered with leaflets. Morphological analysis of these galls suggest an organised spatial distribution of the primordia that are covered with leaflets, both according to the fundamental rules of phyllotaxis.

Based on the experimental information provided here it became possible to identify a complex of factors that can be used to induce fasciation independently from the bacteria as a whole. This complex of factors comprises at least the fas products and/or the att products and in addition other gene products encoded by the pFiD188 plasmid.

09380406-134495

The unique property of R. fascians to amplify shoot primordium formation through the modulation of specific hormone balances in a broad spectrum of plants is useful for the establishment of a regeneration program for a diverse range of plants as will be illustrated in the following examples. Infection of plants with R. fascians leads to the local amplification of shoot primordia, combined with a suppression of further shoot growth and development. Upon amplification, primordia can be released from the suppressive effects of R. fascians by killing of the bacteria. This can be achieved by detaching galls and growing them on media with doses of bactericides like penicillin's. Any other treatment of the galls that will kill the inciting bacteria without harming the plant tissue can be envisioned here.

As a result, from the gall tissues shoots will sprout that can be detached and that will form roots, ultimately giving rise to normal, regenerated plants. This effect is even more dramatic when the galls are treated in the dark, where the release of shoots is accompanied by etiolation, producing masses of elongated sprouts that can be propagated to a normal plant.

This technique has been successfully applied to leafy galls on Nicotiana tabacum, Papaver somniferum, Artemisia annua, Vigna unguiculata, Atropa belladonna, Sesbania rostrata and Digitalis lanata.

The Figure is an overview showing the steps required for micropropagation using either classical procedure or Rhodococcus fascians.

This invention as described here will be further illustrated in the following examples.

654247" 90708660

EXAMPLES**EXAMPLE 1**

Regeneration of plants from leafy galls of various plant species-induced by Rhodococcus fascians

5

1. General method

The following method is applicable to plant species in general. In these examples the plant species Nicotiana tabacum, Papaver somniferum, Artemisia annua,
10 Vigna unguiculata, Atropa belladonna, Sesbania rostrata and Digitalis lanata were used.

1.1. Bacterial strain

Rhodococcus fascians strain D188 or other
15 virulent strains are cultured in solid or liquid conventional bacteria medium or a recommended one containing: Glucose (10 g/l), Yeast extract (5 g/l), Peptone (5 g/l), Agar (15 g/l), distilled water 1l, the pH adjusted to 7.0. The cultures are then incubated in
20 the dark at 25°C.

1.2. Plant material

Selected plant species are used for the propagation purposes. After surface disinfection of the
25 explants (for example: cutting consisting of stem internode fragments with axillary buds) using appropriate reagents known in the art, the explant is transferred to in vitro culture condition using conventional tissue culture medium such as the well-known basal Murashige and
30 Skoog medium (Physiol. Plant. 15, 473-497) or Gamborg medium (Gamborg, O.L. et al., Ex. Cell. Res. 50, 151 (1986)).

1.3. Infection procedure

35 The axenic explant is directly infected with the fast growing bacteria or infected under vacuum. All other method of infection which make close contact of the bacteria and the plant tissue or cells are also suitable.

09380406-12495

WO 98/36635

PCT/EP98/01117

14

In case of Sesbania rostrata, cotyledonous explants were used for direct infection with Rhodococcus fascians. Preferably wounded cotyledons were used for this infection in order to obtain a more efficient infection.

5

1.4. Incubation of the infected explant

Containers containing the infected plant material are incubated in the room culture at 23°C. Depending on the plant species, leafy galls came to be visible after 3-4 weeks of incubation.

2. Regeneration of plants Leafy galls structure

2.1. General procedure

The leafy galls are first isolated from the original explants. Then they are subcultured in a hormone-free conventional plant tissue culture medium additionated with bactericide reagent such as Claforan at final concentration of 500 mg/l. The cultures are preferably incubated in the dark at appropriate temperature (depending on the plant species, for example 19°C for Papaveraceae species; 23°C for Solanaceae species).

2.2. Results

The dark condition combined with the antibiotic treatment improve the shoot elongation. The etiolated shoots thus obtained are then subcultured in a hormone-free culture medium and incubated in light condition. The light intensity as well as the temperature of incubation are selected according to the specific requirement of each plant species under investigation. After several days of cultures, spontaneous root appeared, and thus plantlet are recovered.

Using the described method and in the case of Digitalis lanata, an average of 60 new regenerated shoots/explant are obtained instead of 12 new regenerated shoots using the classical micropropagation method (using exogenous cytokinins).

0930103-1449

WO 98/36635

PCT/EP98/01117

15

Using common method for acclimatization, these new regenerated plantlets are successfully transferred to the green house.

Quantitative and qualitative analysis of
5 cardenolides in the new regenerated plants showed a distribution quite similar to that of the mother plant. In addition the morphological traits of these new regenerated plants are similar to those of the mother plant, indicating a reduced variability as compared to
10 the classical micropropagation technique.

EXAMPLE 2

Germplasm storage

1. Short-term conservation

15 The leafy galls containing the bacteria are isolated from the original explant then subcultured in a conventional plant tissue culture medium with reduced sugar concentration (15g/l of sucrose). After one week of culture at 20°C, the culture containers are transferred
20 to 4°C under dark condition. In such conditions the capability of the shoots to develop (after bactericide treatment) is maintained for 1-2 years.

2. Long term conservation

25 The well established procedure for cryo-conservation using liquid nitrogen is also convenient for the leafy gall structure (George E.F., in: Plant Propagation by Tissue Culture, Part I, George Edwin F. (Ed.), Exegetics Ltd, Edington, Wilts, England, pp. 163-
30 181 (1993)).

054627-1-1490

PCT/EP98/01117

16

CLAIMS

1. Method for the in vitro preparation of starting materials for micropropagation and/or storage of germplasm, comprising the steps of:

- a) contacting plant material with a micro-
5 organism that induces fasciation and/or one or more fasciation-inducing factors derived from the micro-organism or derived from the infected and/or fasciated tissue;
- b) developing leafy galls or shoot outgrows on
10 the plant material; and
- c) isolating the leafy galls or shoot outgrows as the starting materials.

2. Method for the in vitro micropropagation of plants, which method comprises the method as claimed in
15 claim 1, further characterized by the steps of

- d) eliminating or inactivating the micro-organism that induces fasciation and/or the one or more fasciation-related factors derived from the micro-organism or derived from the infected and/or fasciated
20 tissue;
- e) culturing the leafy gall or shoot outgrows in or on one or more suitable culture media to allow shoot and root development for obtaining plantlets; and
- f) transferring the plantlets thus obtained,
25 optionally after acclimatisation, to conventional growing conditions to obtain regenerated plants.

3. Method for the preservation of plants or germplasm, which method comprises the method as claimed in claim 1, further characterized by the step of storing
30 the leafy galls or shoot outgrows under growth limiting conditions.

4. Method as claimed in claims 1-3, wherein the fasciation-inducing micro-organism is Rhodococcus fascians.

35 5. Method as claimed in claim 3 and 4, wherein the growth limiting conditions comprise low temperatures

0330406-4499

WO 98/36635

PCT/EP98/01117

17

of about 4°C or cryo-conserving conditions, such as storage in liquid nitrogen.

6. Method as claimed in claims 1-5, wherein the plant material comprises explants, such as leaf discs, 5 organs, tissue fragments, seedlings, seeds, embryos, isolated cells, protoplasts, cell cultures and callus tissues.


7. Method as claimed in claims 1-6, wherein the plant material is material of species of the genera 10 Brassicaceae, Salicaceae, Compositae, Solanaceae, Scrophulariaceae, Liliaceae, Papaveraceae, Gramineae and Fabaceae.

8. Method for the in vitro micropropagation of plants, which method comprises the method of claim 3 and 15 optionally the subsequent removal or inactivation of still present micro-organisms that induce fasciation and/or fasciation related factors derived from the micro-organism or derived from the infected and/or fasciated tissue, culturing the leafy gall or shoot outgrows in or 20 on one or more suitable culture media to induce shoot and root development for obtaining plantlets, and transferring the plantlets thus obtained, optionally after acclimisation, to conventional growing conditions to obtain regenerated plants.


09380105-112490

6642T" 90T0860

Classical procedure using in vitro tissue culture method



Use of Rhodococcus fascians for micropropagating plants



WO 98/36635

PRINT TIME AUG.18. 10:33AM


RECEIVED TIME AUG.18. 10:01AM

09/380106

PCT/EP98/01117

Page 1

Attorney's Docket No. DECLE2.001AFC

**DECLARATION - USA PATENT APPLICATION**

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name;

I believe I am an original, first and joint inventor of the subject matter which is claimed and for which a patent is sought on the invention entitled **PLANT MICROPROPAGATION AND GERMPLASM STORAGE**; the specification of which was filed as International Application No. PCT/EP98/01117 on 19 February 1998;

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above;

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, § 1.56;

I hereby claim foreign priority benefits under Title 35, United States Code, § 119(a)-(d) of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

PRIOR FOREIGN APPLICATION(S)**Priority
Claimed**

No.: 97200513.6
97200701.7

Country: Europe
Europe

Date Filed: 21 February 1997
10 March 1997

Yes
Yes

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful, false statements may jeopardize the validity of the application or any patent issued thereon.

Full name of first inventor: Koenraad Hilkuire Maria Goethals

Inventor's signature [Signature]


Date 25-08-99

Residence: Bollewerkstraat 85, B-9031 Gent, Belgium BE

Citizenship: Belgium

Page 2

Attorney's Docket No. DECLE2.001APC

2.00
Full name of Second inventor: Mondher El JaziriInventor's signature Date 01-09-99Residence: Place de la Gare 10, B-1150 Woluwe Saint Pierre, Belgium BELCitizenship: BelgiumPost Office Address: same as above3.00
Full name of Third inventor: Marc Charles Ernest Van MontaguInventor's signature Date 20 / VII / 99Residence: De Stassartstraat 120, B-1050 Brussels, Belgium BELCitizenship: BelgiumPost Office Address: same as above

Send Correspondence To:

KNOBBE, MARTENS, OLSON & BEAR, LLPCustomer No. 20,995E:\DOCS\DEA\DEA-1567.DOC:kac
081099